

NANORMEN® Oral suspension

NEMATICIDAL DEWORMER



Nanormen® Oral Suspension, is indicated for the treatment and control of gastrointestinal nematodes in dogs and cats of any age.

Nanormen® is effective against *Toxocara canis*, *Toxocara cati*, *Toxascaris leonina*, *Ancylostoma caninum*, *Ancylostoma tubaeforme* and *Uncinaria stenocephala*.

Technical Specification

SPECIES

Dogs and cats.

DOSAGE FORM

Oral suspension

THERAPEUTIC ACTION

Nematicidal dewormer

COMPOSITION

Each mL contains:

Pirantel Pamoate 144.13 mg

(Equivalent to 50 mg of Pirantel base)

Excipients q.s.p 1 mL

INDICATIONS

Nanormen® Oral Suspension, is indicated for the treatment and control of gastrointestinal nematodes in dogs and cats of any age.

Nanormen® is effective against *Toxocara canis*, *Toxocara cati*, *Toxascaris leonina*, *Ancylostoma caninum*, *Ancylostoma tubaeforme* and *Uncinaria stenocephala*.

ROUTE OF ADMINISTRATION AND DOSAGE

Administration way: Oral.

Dose of the active substance:

- Dogs: 5 mg / Kg of weight.
- Cats: 20 mg / Kg of weight.

Product dosage:

- **Dogs: administer 2 drops per Kg of weight or 1 mL for every 10 Kg of weight in a single dose.**
- **Cats: Administer 8 drops per Kg of weight or 2 mL for every 5 Kg of weight in a single dose.**

DRUG INTERACTIONS

Pirantel can interact with medications containing Piperazine, Levamisole, or other Nicotine-like cholinergic drugs.

CONTRAINDICATIONS

- Do not administer to severely weakened animals.
- Do not administer to pregnant or lactating females.
- Do not administer to hypersensitive animals to Pirantel Pamoato.

SPECIAL PRECAUTIONS FOR USE

Special warnings and precautions for use:

- Shake before using.
- Keep out of the reach of children.
- Nanormen® is recommended to treat and control parasite reinfestation in puppies and kittens, for which animals can be treated at 2, 4, 6, 8 and 10 weeks of life.
- Animals exposed to highly contaminated environments, it is recommended to administer Nanormen® every 2 or 3 months to control reinfestation by *Toxocara canis*.

SPECIAL PRECAUTIONS FOR THE OPERATOR

Do not handle by people hypersensitive to Pirantel Pamoato

SIDE EFFECTS

Mild and transient side effects may occasionally occur, including gastrointestinal signs such as nausea, vomiting, anorexia, abdominal pain, diarrhea, erythema of the skin, and increased liver enzyme values.

ADVERSE EFFECTS

Mild and transient side effects may occasionally occur, including gastrointestinal signs such as nausea, vomiting, anorexia, abdominal pain, diarrhea, erythema of the skin, and increased liver enzyme values.

OBSERVATIONS

Special precautions for the disposal of unused product or waste material:

Empty containers can be discarded as household waste, without any special precautions. Do not dispose of containers with product residues on the ground or water courses. For expired or unused products, contact the manufacturing laboratory.

CONSERVATION

Store between 2 ° and 30 ° C, protected from light. Once opened, use within 12 weeks and discard the excess product.

CONDITION OF SALE

sale with Veterinary Medical prescription

PRESENTATION

20 mL bottle

PREPARED BY

Laboratorio Drag Pharma Chile Invetec S.A.

RECORDS

Chile: Reg. SAG N° 178

Costa Rica: Reg. N° MAG CL4-42-10-5936

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian