

# PANVERMIC® LARGE BREED - Oral Tablet

## BROAD SPECTRUM INTERNAL ANTIPARASITIC.



Broad spectrum internal antiparasitic in a single dose. Effective against nematodes and cestodes, such as: *Toxocara canis*, *Toxascaris leonina*, *Ancylostoma caninum*, *Hookworm stenocephala*, *Taenia hydatigena*, *Taenia pisiformis*, *Taenia taeniformis*, *Dipylidium caninum*, and *Echinococcus granulosus*.

## Technical Specification

### SPECIES

Large breed dogs.

### DOSAGE FORM

Oral Tablet

### THERAPEUTIC ACTION

Broad spectrum internal antiparasitic.

### COMPOSITION

Each tablet contains:

Praziquantel...	175 mg
Mebendazole ...	770 mg
Pyrantel Pamoate ...	.504 mg
(Equivalent to 175 mg of Pyrantel base)	
Excipients q.s.p	1 tablet

### INDICATIONS

Broad spectrum internal antiparasitic in a single dose. Effective against nematodes and cestodes, such as: *Toxocara canis*, *Toxascaris leonina*, *Ancylostoma caninum*, *Hookworm stenocephala*, *Taenia hydatigena*, *Taenia pisiformis*, *Taenia taeniformis*, *Dipylidium caninum*, and *Echinococcus granulosus*.

### ROUTE OF ADMINISTRATION AND DOSAGE

Administer orally, directly into the oral cavity or mixed with food.

**- Dose of active ingredients:**

Administer 22 mg / Kg of Mebendazole; 5 mg / Kg of Praziquantel and 5 mg / Kg of Pyrantel as a base.

**- Product dosage:**

1 tablet for every 35 kg of weight. Single dose.

For the treatment of *Trichuris vulpis* infestations, administer 1 tablet for every 35 kg of weight for 3 consecutive days.

#### CONTRAINDICATIONS

---

- Do not administer to dogs with liver failure, malnourished, dehydrated or anemic.
- Do not administer in dogs less than four weeks of age.
- Interactions with other pharmaceuticals:
- Do not administer together with Piperazine as it could antagonize the effect of Pyrantel Pamoate.
- Do not administer in conjunction with Morantel or Levamisole due to their mechanisms of action and toxicity similar to Pyrantel.

#### SPECIAL PRECAUTIONS FOR THE OPERATOR

---

Wash your hands after administering the product.

#### WARNINGS

---

Keep out of the reach of children.

#### ADVERSE EFFECTS

---

- In some animals loose stools and in some cases moderate diarrhea may occur.
- In less than 5% of treated dogs there may be anorexia, vomiting and lethargy.

#### CONSERVATION

---

- Keep in a dry place and protected from light.
- Store at room temperature between 15 and 30 ° C

#### CONDITION OF SALE

---

Sale with veterinary prescription.

#### PRESENTATION

---

Box with 1 tablets

#### PREPARED BY

---

Laboratorio Drag Pharma Chile Invetec S.A.

#### RECORDS

---

**Chile:** Reg. SAG N° 2076

**Bolivia:** Reg. SENASAG PUV-F-N° 005508/13

#### COUNTRIES WHERE IT IS MARKETING

---

Imported and distributed in Bolivia by:

ZOOFARMA  
TEL: +(591)222-3357  
Street Díaz Romero 1339, La Paz.

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian