

SINPULSPOT[®] PLUS Large Breed - Spot on

EXTERNAL ANTIPARASITIC. SPOT-ON



Effective insecticide in the control and prevention of infestation by fleas (*Ctenocephalides felis*) and ticks (*Rhipicephalus sanguineus*) in dogs. The product provides effective protection against reinfestations for 4 weeks after application.

Technical Specification

SPECIES

Dogs over 15 Kg of body weight.

DOSAGE FORM

Topical solution. Spot on

THERAPEUTIC ACTION

External antiparasitic. Spot-on

COMPOSITION

Each dose of 2.0 mL contains:
Permethrin (cis/trans 25/75)..... 650 mg
Piperonyl Butoxide.....650 mg
Methoprene.....75 mg
Excipients q.s.....2.0 mL

INDICATIONS

Effective insecticide in the control and prevention of infestation by fleas (*Ctenocephalides felis*) and ticks (*Rhipicephalus sanguineus*) in dogs. The product provides effective protection against reinfestations for 4 weeks after application.

MODE OF APPLICATION

Cut off the tip of the pipette (s). Separate the hair in the area of application, so that the dose is administered directly to the skin.
- In dogs over 15 kg apply a 2 mL pipette at the withers and another 2 mL pipette at the base of the tail (SINPULSPOT[®] PLUS, Large Breed).

ROUTE OF ADMINISTRATION AND DOSAGE

Dose: 4 mL (2 pipettes).

Single dose. Apply the recommended dose every 4 weeks.

DRUG INTERACTIONS

Do not use concomitantly with other pharmaceuticals.

CONTRAINDICATIONS

- Do not apply to dogs younger than 4 months.
- **Do not apply to cats.**
- Do not apply to pregnant or lactating females.

SPECIAL PRECAUTIONS FOR USE

Special warnings and precautions for use:

- Keep out of the reach of children.
- For a greater residual effect, it is recommended to apply SINGULSPOT® PLUS, 3 days after the last bath of your dog, without re-bathing until it is strictly necessary.

SPECIAL PRECAUTIONS FOR THE OPERATOR

- Avoid contact with skin: wash your hands after handling the product.
- Avoid contact with eyes and in case of splashes wash with plenty of water.

ADVERSE EFFECTS

Unwanted effects and adverse reactions:

On very rare occasions, slight coloration may occur at the application site in light-coated dogs; transient signs of skin irritation such as redness, erythema, or itching; behavioral changes such as restlessness; and / or gastrointestinal signs such as salivation, vomiting or loss of appetite, which should disappear 24 hours after application. If signs of irritation persist, it is recommended to bathe the dog with a mild shampoo to remove the product and rinse with plenty of water. If the signs described continue or worsen, consult a Veterinary Doctor.

OBSERVATIONS

External use only.

Poison

Special precautions for the disposal of unused product or waste material:

Discard the remains of unused product in its original container. Do not discard the empty container or with product residues in rivers, lakes or streams of natural water. The product is toxic to aquatic organisms. Dispose of this product with caution with household waste.

CONSERVATION

Keep in a fresh, dry place, at room temperature between 2 and 30 °C.

CONDITION OF SALE

OTC product (non-prescription).

PRESENTATION

2 pipettes of 2 mL

PREPARED BY

Drag Pharma Laboratory.

RECORDS

Chile: Reg. SAG N°: 1672

Costa Rica: Reg. N° MAG CL4-43-04-4139

COUNTRIES WHERE IT IS MARKETED

Imported and distributed in Bolivia by:

ZOOFARMA

TEL: +(591)222-3357

Street Díaz Romero 1339, La Paz.

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian