

SINPULKILL® Dog Collar

EXTERNAL ANTIPARASITICS.



Effective antiparasitic collar in the treatment and control of infestation by fleas (*Ctenocephalides felis*) and ticks (*Rhipicephalus sanguineus*) in dogs, guaranteeing flea control for 4 months, and ticks for 3 months.

Technical Specification

SPECIES

Dogs

DOSAGE FORM

Collar.

THERAPEUTIC ACTION

External Antiparasitics.

COMPOSITION

Each 100 g of collar contains:

Diazinon 15.0 g

Excipients q.s.p 100.0 g

INDICATIONS

Effective antiparasitic collar in the treatment and control of infestation by fleas (*Ctenocephalides felis*) and ticks (*Rhipicephalus sanguineus*) in dogs, guaranteeing flea control for 4 months, and ticks for 3 months.

USE INSTRUCTIONS

Attach the collar loosely to the dog's neck and cut off the remaining collar protruding from the buckle. It is recommended to leave a space of two fingers between the collar and the neck.

DRUG INTERACTIONS

- It is recommended not to use organophosphates together with other cholinesterase inhibitor compounds, especially antiparasitics that use such a mechanism of action.

- Do not use concomitantly with other antiparasitic products.

CONTRAINDICATIONS

- Do not use in sick or convalescent animals.
- Do not use in pregnant, lactating or breeding animals.
- Do not use in dogs under 6 months of age.
- Do not administer to animals hypersensitive to organophosphate compounds, such as Diazinon, preferably Greyhound and Whippet breeds.

SPECIAL PRECAUTIONS FOR USE

- Prevent the animal from playing, licking or swallowing the collar.
- Remove the collar in case of irritation in the contact area, or if signs such as salivation, vomiting or diarrhea appear.
- For external use only.
- Keep out of the reach of children.

SPECIAL PRECAUTIONS FOR THE OPERATOR

- Do not handle by people who are hypersensitive to organophosphate compounds, such as Diazinon.
- Do not handle by pregnant or lactating women.
- Wear gloves when handling the collar, either when putting it on or taking it off.
- Do not smoke, eat or drink near the collar.
- In case of accidental ingestion, go immediately to a medical center and show the product label.
- Prevent the owner or children from sleeping with the animal while it is wearing the collar.
- Wash your hands after handling the collar.
- Prevent children from playing with the collar.

OBSERVATIONS

It not contains metallic parts for preventing allergies.

Special precautions for disposal of unused product or waste material:

- Empty containers can be disposed of as household waste, without any special precautions.
- Do not dispose of containers with product remains on the ground or water courses.
- For expired or unused products, contact the manufacturing laboratory.

CONSERVATION

Store between 15 and 30 ° C, protected from light.

CONDITION OF SALE

OTC product (non-prescription).

PRESENTATION

Case with 1 Collar

PREPARED BY

Manufactured by: Societé AB7 Industries S.A. Chemin des Monges Deyme, 314150 Montgiscard.

France.

Exported by:

Vetpharma Animal Health S.L. Les Corts, 23. 08028 Barcelona. Spain.

Imported and distributed in Chile by Laboratorio Drag Pharma Chile Invetec S.A.

RECORDS

Reg. S.A.G. No.: 1646

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian