

# TOPFLAM<sup>®</sup> Topical Gel

---

## ANTI-INFLAMMATORY



Topflam<sup>®</sup> is indicated for the control of pain and inflammation associated with the musculoskeletal system.

## Technical Specification

### SPECIES

---

Horses.

### DOSAGE FORM

---

Topical Gel

### THERAPEUTIC ACTION

---

Anti-inflammatory

### COMPOSITION

---

Each 100 g of product contains:  
Diclofenac Sodium ..... 1 g  
Excipients q.s.p ..... 100 g

### INDICATIONS

---

Topflam<sup>®</sup> is indicated for the control of pain and inflammation associated with the musculoskeletal system.

### ROUTE OF ADMINISTRATION AND DOSAGE

---

Administer topically 7.3 g of gel or a 13 cm band of gel, twice a day for 5 days, equivalent to 73 mg of Diclofenac Sodium in each application. Rub the joint to be treated with the product until it is completely absorbed.

### DRUG INTERACTIONS

---

The use of other topical drugs should be avoided as they can inhibit the absorption of Topflam<sup>®</sup> and

cause irritation in the area of application. The concomitant administration of other NSAIDs by the systemic route has not been studied.

#### CONTRAINDICATIONS

---

- Do not administer in animals sensitive to the drug.
- Do not use in conjunction with NSAIDs.
- Do not administer during pregnancy or lactation.
- Do not administer to foals under one year of age.

#### SPECIAL PRECAUTIONS FOR USE

---

Exceeding the recommended dose or treating several joints at the same time in the same animal, can increase the plasma concentrations of Diclofenac.

Keep out of reach of children and pets.

#### SPECIAL PRECAUTIONS FOR THE OPERATOR

---

- Avoid contact with the skin, wear gloves to apply the product. In the case of contact, wash with soap and water.
- Avoid accidental contact with the eyes. In the case of contact, rinse with plenty of water and consult a doctor immediately.
- Do not handle by people who are hypersensitive to Diclofenac Sodium.
- In case of accidental ingestion, go immediately to a medical center and show the product label.

#### WARNINGS

---

Keep out of the reach of children.

#### GUARD PERIOD

---

Do not administer to horses whose meat is intended for human consumption.

#### OBSERVATIONS

---

##### **Special precautions for disposal of unused product or waste material:**

Empty containers can be disposed of as household waste, without any special precautions. Do not dispose of containers with product remains on the ground or water courses. For expired or unused products, contact the manufacturing laboratory.

#### CONSERVATION

---

Keep in a cool and dry place, at room temperature between 15° and 30° C, protected from light. Once the container is opened, use within 12 weeks. Discard unused product after that time period.

#### CONDITION OF SALE

---

Sale with veterinary prescription.

#### PRESENTATION

---

120 g tube

#### PREPARED BY

---

Laboratorio Drag Pharma Chile Invetec S.A.

#### RECORDS

---

**Chile:** Reg. SAG N° 2042

**Perú:** Reg. SENASA F.C4.75.I.0010

#### COUNTRIES WHERE IT IS MARKETED

---

Imported and Distributed in Peru by Representaciones Durand SAC.  
Av. Manuel Olgún No. 501 Office No. 604 Santiago de Surco Lima.

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian