

TRANSIMED® Otic and Topical Suspension

ANTIBACTERIAL, ANTIFUNGAL AND STEROIDAL ANTI-INFLAMMATORY



Transimed® is indicated for the treatment of external otitis and dermatitis caused by fungi, yeasts and Gram (-) bacteria.

Technical Specification

SPECIES

Dogs and cats.

DOSAGE FORM

Otic and topical suspension.

THERAPEUTIC ACTION

Antibacterial, antifungal and steroidal anti-inflammatory

COMPOSITION

Each 1 mL contains:

Miconazole Nitrate 23.0 mg
Prednisolone Acetate 5.0 mg
Polymyxin B Sulfate 5,000 IU
Excipients q.s.p 1 mL

INDICATIONS

Transimed® is indicated for the treatment of external otitis and dermatitis caused by fungi, yeasts and Gram (-) bacteria.

ROUTE OF ADMINISTRATION AND DOSAGE

Route of administration: Topical.

Dosage and directions for use:

In otitis:

- Clean the external auditory canal, with a ceruminolytic and antiseptic product.
- Instill 5 drops of Transimed® in the external auditory canal 2 to 3 times a day, for up to 15 days.

In dermatitis:

- Cut the hair in the affected area and clean it.
- Spread several drops of Transimed® on the area to be treated, 2 to 3 times a day for up to 10 days.

DRUG INTERACTIONS

- Polymyxin B has interactions with aminoglycoside antibiotics, Bacitracin, Quinidine, Quinine.
- Sodium Citrate enhances the nephrotoxic and neurotoxic potential of Polymyxin B.
- Succinylcholine and Tubocurarine can prolong the neuromuscular blockade of respiratory paralysis associated with the use of Polymyxin B.

CONTRAINDICATIONS

- Do not use in animals with a perforated eardrum due to the ototoxic effect of Polymyxin B.
- Do not use in pregnant or lactating females.

SPECIAL PRECAUTIONS FOR USE

In case of prolonged therapy (more than 7 days) it should be gradually suspended in successive days according to what is indicated by the treating Veterinarian.

SPECIAL PRECAUTIONS FOR THE OPERATOR

- Do not handle by people who are hypersensitive to glucocorticoids.
- In case of contact with skin, eyes or mucous membranes, wash immediately with plenty of water.
- Do not handle by pregnant women.
- In case of ingestion, go immediately to a medical center and show the product label.

WARNINGS

Keep out of the reach of children.

ADVERSE EFFECTS

- The use of Transimed® for more than 7 days could delay wound healing.
- The use of Transimed® may be associated with the development of deafness or partial hearing loss in a small number of sensitive (geriatric) dogs. The hearing deficit is usually temporary. If vestibular or hearing dysfunctions are detected during the course of treatment, therapy should be discontinued and the external auditory canal should be flushed.

OBSERVATIONS

Shake before using.

Special precautions for disposal of unused product or waste material:

Empty containers can be disposed of as household waste, without any special precautions. Do not dispose of containers with product remains on the ground or water courses. For expired or unused products, contact the manufacturing laboratory.

CONSERVATION

Store at room temperature, between 15 and 30 ° C. Once opened use the product within 3 weeks. Discard the unused product after that period of time.

CONDITION OF SALE

Sale with veterinary prescription.

PRESENTATION

15 mL bottle with cannula.

PREPARED BY

Laboratorio Drag Pharma Chile Invetec S.A.

RECORDS

Chile: Reg. SAG N° 0526-B
El Salvador: VE2015105122

COUNTRIES WHERE IT IS MARKETED

Distribution in El Salvador:
Rafael Alfredo Alfaro Castillo.
8th C. Pte. Pje. Moreno N ° 112, Col. Flor Blanca.
San Salvador, El Salvador.

Imported and distributed in Bolivia by:
ZOOFARMA
TEL: +(591)222-3357
Street Díaz Romero 1339, La Paz.

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian