



BROAD SPECTRUM ANTIFUNGAL FOR HORSES



Technical Specification

SPECIES

Horses

DOSAGE FORM

Topical solution

THERAPEUTIC ACTION

Broad spectrum antifungal for horses

COMPOSITION

Each 1 mL of topical solution contains:

Enilconazole.....100 mg

Excipients q.s.....1 mL

MODE OF APPLICATION

- Before starting the treatment with the diluted solution of Equinyl®, the horse must be cleaned and/or bathed with a brush and mild soap, removing any scabs and scales that exist to facilitate the penetration of the active ingredient into the coat.
- The first two applications should be carried out covering the entire body of the animal, with the aim of also reaching subclinical lesions and spores present in the fur. The diluted solution should then be applied only to the lesions, covering a large area around the affected skin.

ROUTE OF ADMINISTRATION AND DOSAGE

The Equinyl® solution must be diluted before use, adding 1 part of the solution to 50 parts of warm water until obtaining a 0.2% Enilconazole emulsion (2 mg of active substance / mL of water), which it should be used to treat the horse topically.

It is recommended to follow the following dilution guidelines:

- 10 mL of Equinyl®, concentrated, must be diluted in ½ liter of warm water.
- 20 mL of Equinyl®, concentrated, must be diluted in 1 liter of warm water.
- 100 mL of Equinyl®, concentrated, must be diluted in 5 liters of warm water.
- 120 mL of Equinyl®, concentrated, must be diluted in 6 liters of warm water.

Frequency and duration of treatment

The diluted Equinyl® solution should be applied at intervals of 3 to 4 days. The duration of treatment can vary from a minimum of 4 applications to a maximum of 12 applications in severe cases. Treatment should continue until after clinical resolution is observed and negative cultures are obtained.

DRUG INTERACTIONS

No specific interactions have been reported. However, like other azoles, systemic treatment may result in inhibition of the cytochrome P450 enzyme.

CONTRAINDICATIONS

- Do not administer to animals with known hypersensitivity to Enilconazole.
- Do not use in horses less than 6 months of age.
- Do not wet the animal's mucous membranes (oral and ocular) with the product.

SPECIAL PRECAUTIONS FOR USE

- Keep out of the reach of children and pets.
- The diluted product must be used immediately.

SPECIAL PRECAUTIONS FOR THE OPERATOR

- Use gloves and protective glasses when preparing and applying the product.
- Wash your hands with plenty of soap and water after handling and applying the product.
- Do not smoke, eat or drink while handling and applying the product.
- Apply in ventilated environments.
- Avoid contact with the treated animal until its fur is completely dry.
- Do not handle by people hypersensitive to Enilconazole.
- In case of contact with skin, remove contaminated clothing and wash immediately with plenty of soap and water.
- In case of contact with eyes, wash immediately with plenty of water. In case of irritation, go to a medical center and show the product label.
- In case of accidental ingestion, go immediately to a medical center and show the product label.

WARNINGS

- Keep out of the reach of children and pets.
- The diluted product must be used immediately.

ADVERSE EFFECTS

- The development of adverse effects in animals treated with the recommended dose is unlikely, because absorption through the skin is very low.
- When the Equinyl® product is used as indicated, no adverse effects have been recorded.

GUARD PERIOD

Meat: zero days

CONSERVATION

Store at room temperature no higher than 30°C. Do not refrigerate or freeze. Once the container is opened, use within 8 weeks. Discard unused product after that period of time. Use immediately once prepared and discard any excess dilution.

CONDITION OF SALE

Sale under Veterinary Medical prescription.

PREPARED BY

Prepared and distributed by:
Laboratory Drag Pharma Chile Invetec S.A.
Lautaro No. 300. Quilicura. Santiago. Chili
www.dragpharma.cl

RECORDS

Reg. SAG N° 2586

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian