


CAVIVET® Injectable solution

RESTORATIVE TONIC



Tonic restorative adjuvant in the treatment of various pathologies.

- **Horses:** Failures of development and growth; fertility problems; improvement of skin and coat, improvement in the training and in preparation for rodeo participation.
- **Bovine:** Strengthening of calves, strengthening of cows in the last trimester of gestation; fertility problems; performance failure in bulls; and treatment of milk fever in cattle.
- **Pigs:** Strengthening of piglets, breeding sows and boars in mounting.
- **Ovine:** Strengthening of rams and sheep in the last trimester of gestation.

Technical Specification

SPECIES

Bovines, horses, pigs and sheep.

DOSAGE FORM

Injectable solution.

THERAPEUTIC ACTION

Restorative tonic

COMPOSITION

Each 100 mL of injectable solution contains:

Calcium Gluconate Monohydrate.....	5.0000 g
Calcium D-Saccharate Tetrahydrate.....	0.0800 g
Calcium Levulinate Dihydrate.....	1.0000 g
Calcium Glycerophosphate Anhydrous.....	0.5000 g
Magnesium Glycerophosphate Dihydrate.....	1.5000 g
Sodium Glycerophosphate x 5.5 H ₂ O.....	0.8000 g
Sodium selenite Pentahydrate.....	0.0333 g
Cobalt Chloride Hexahydrate.....	0.0800 g
Potassium Chloride.....	0.4000 g
Excipients q.s.p.....	100 mL

INDICATIONS

Tonic restorative adjuvant in the treatment of various pathologies.

- **Horses:** Failures of development and growth; fertility problems; improvement of skin and coat, improvement in the training and in preparation for rodeo participation.
- **Bovine:** Strengthening of calves, strengthening of cows in the last trimester of gestation; fertility problems; performance failure in bulls; and treatment of milk fever in cattle.
- **Pigs:** Strengthening of piglets, breeding sows and boars in mounting.
- **Ovine:** Strengthening of rams and sheep in the last trimester of gestation.

ROUTE OF ADMINISTRATION AND DOSAGE

Administration route:

Intramuscular, subcutaneous, slow intravenous.

Dose:

- **Cattle and horses:** 15 - 20 mL in a single dose. The treatment can be repeated after one week.
- **Sheep and pigs:** 10 mL in a single dose. The treatment can be repeated after one week.

CONTRAINDICATIONS

Do not administer simultaneously with Tetracyclines.

WARNINGS

Special warnings and precautions for use:

- Endovenous administration could cause phlebitis problems.
- Endovenous administration should be done slowly due to the risk of bradycardia or cardiac arrest.
- Keep out of the reach of children and pets.

GUARD PERIOD

Meat: 0 days.

Milk: 0 days.

OBSERVATIONS

Special precautions for the disposal of unused product or waste material:

Empty containers can be discarded as household waste, without any special precautions. Do not dispose of containers with product residues on the ground or water courses. For expired or unused products contact the manufacturer laboratory.

CONSERVATION

Store between 15 ° and 30 ° C, protected from light.

CONDITION OF SALE

To be supplied only on veterinary prescription.

PRESENTATION

250 mL ampoule bottle

PREPARED BY

Drag Pharma Laboratory.

RECORDS

- Chile: Reg. SAG No. 937
- Costa Rica: Reg. No. MAG CL4-32-02-3648

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian