

# DIARREVET® Oral Powder

---

## ANTI-DIARRHEAL.



Control diarrhea caused by germs sensitive to sulfonamides (*Salmonella spp*, *E.coli* and *Coccidia*).

## Technical Specification

### SPECIES

---

Bovine, horses, pigs, ovine and goats.

### DOSAGE FORM

---

Oral powder.

### THERAPEUTIC ACTION

---

Anti-diarrheal.

### COMPOSITION

---

Each 100 g contains:

Sulfaguanidine Monohydrate .....	15 g
Phthalylsulfathiazole .....	5 g
Citrus Pectin .....	1 g
Bentonite .....	2 g
Colloidal Kaolin .....	77 g

### INDICATIONS

---

Control diarrhea caused by germs sensitive to sulfonamides (*Salmonella spp*, *E.coli* and *Coccidia*).

### ROUTE OF ADMINISTRATION AND DOSAGE

---

Oral route:

Dissolve the recommended dose in 100 to 200 mL of warm water.

**Product dose:** (1 tablespoon equal to 10 g of the product)

	Bovine and adult horses	Ovines, Goats, Calves and Fools	Adult Pigs	Piglets
<b>INITIAL DOSE</b>	1 tablespoon (for every 500 Kg of weight: Sulfaguanidine 3g; Phthalysulfatiazole 1g; Pectin 0,2g; Bentonite 0,4g; Kaolin 15,4g)	1 tablespoon (for every 50 Kg of weight: Sulfaguanidine 1,5g; Phthalysulfatiazole 0,5g; Pectin 0,1g; Bentonite 0,2g; Kaolin 7,7g)	1 ½ tablespoon (for every 100 Kg of weight: Sulfaguanidine 2,25 g; Phthalysulfatiazole 0,75 g; Pectin 0,15 g; Bentonite 0,3 g; Kaolin 11,55 g)	½ tablespoon (for every 30 Kg of weight: Sulfaguanidine 0,75 g; Phthalysulfatiazole 0,25 g; Pectin 0,05 g; Bentonite 0,1 g; Kaolin 3,85 g)
<b>MAINTENANCE DOSE</b>	1 tablespoon (for every 500 Kg of weight: Sulfaguanidine 1,5 g; Phthalysulfatiazole 0,5 g; Pectin 0,1 g; Bentonite 0,2 g; Kaolin 7,7 g) every 8 hours for 4 days.	1 tablespoon (for every 50 Kg of weight: Sulfaguanidine 1,5 g; Phthalysulfatiazole 0,5 g; Pectin 0,1 g; Bentonite 0,2 g; Kaolin 7,7 g) every 8 hours for 4 days.	1 tablespoon (for every 100 Kg of weight: Sulfaguanidine 1,5 g; Phthalysulfatiazole 0,5 g; Pectin 0,1 g; Bentonite 0,2 g; Kaolin 7,7 g) every 8 hours for 4 days.	¼ tablespoon (for every 30 Kg of weight: Sulfaguanidine 0,375 g; Phthalysulfatiazole 0,125 g; Pectin 0,025 g; Bentonite 0,05 g; Kaolin 1,925 g) every 8 hours for 4 days.

#### DRUG INTERACTIONS

Adsorbents such as Kaolin and Pectin can inhibit the intestinal absorption of other drugs that are administered together or during treatment with DIARREVET®.

#### CONTRAINDICATIONS

Do not use in animals hypersensitive to sulfonamides.

#### SPECIAL PRECAUTIONS FOR USE

Use with caution in animals with kidney or liver failure.

#### WARNINGS

Keep out of reach of children.

#### GUARD PERIOD

**Meat:** 10 days (cattle, pigs, sheep and goats). Do not administer in horses whose meat is intended for human consumption.

**Milk:** 4 days (cattle). Do not administer in sheep or goats whose milk is intended for human consumption.

#### OBSERVATIONS

#### Special precautions for disposal of unused product or waste material:

Empty containers can be disposed of as household waste, without any special precautions. Do not dispose of containers with product remains on the ground or water courses. For expired or unused products, contact the manufacturing laboratory.

#### CONSERVATION

Store between 15 ° and 30 ° C, protected from light. Use immediately once prepared and discard the excess product.

#### CONDITION OF SALE

---

Supply only on veterinary prescription.

#### PRESENTATION

---

Box with 25 sachets of 50 g each.

#### PREPARED BY

---

Drag Pharma Laboratory.

#### RECORDS

---

- Chile: Reg. SAG No. 942
- Dominican Republic: Reg. No. 5608

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian