

# GLICEFAR<sup>®</sup> Injectable solution

---

## CENTRAL MUSCLE RELAXANT



Adjuvant for the induction of anesthesia in horses.

It can be used as an inducing agent in combination with barbiturates to soften the stage of induction and recovery of the latter.

It can also be used in a combination with other agents such as Xylazine and Ketamine, to maintain anesthesia and thus decrease the concentration of inhalation anesthesia.

Its use is recommended only in adult horses.

## Technical Specification

### SPECIES

---

Horses.

### DOSAGE FORM

---

Injectable solution.

### THERAPEUTIC ACTION

---

Central muscle relaxant

### COMPOSITION

---

Each mL of product contains:

Guaifenesin.....100 mg

Excipients q.s.p.....1 mL

### INDICATIONS

---

Adjuvant for the induction of anesthesia in horses.

It can be used as an inducing agent in combination with barbiturates to soften the stage of induction and recovery of the latter.

It can also be used in a combination with other agents such as Xylazine and Ketamine, to maintain anesthesia and thus decrease the concentration of inhalation anesthesia.

Its use is recommended only in adult horses.

### USE INSTRUCTIONS

---

- To use in combination with barbiturates, add the barbiturate (example: Thiopental) dose to the Glicefar<sup>®</sup> solution before the administration and proceed to the infusion or administer it as a bolus immediately after the administration of Glicefar<sup>®</sup>.
- To use in combination with xylazine and/or ketamine, Xylazine is administered 2 to 5 minutes

before the administration of GliceFar® in the recommended dose and later the ketamine will be administered, once the ataxia signs from GliceFar® are present.

#### ROUTE OF ADMINISTRATION AND DOSAGE

---

Intravenous administration only.

**Recommended dose:** 100 - 110 mg / Kg

**Recommended practical dose:** 1 mL / Kg of weight.

#### CONTRAINDICATIONS

---

Do not administer to pregnant or lactating females.

#### SPECIAL PRECAUTIONS FOR USE

---

- The extravascular injection should be avoided as it can cause irritation of the tissue.
- It is recommended to monitor the heart and breathing rate of the patient when administering Guaifenesin (Glyceryl Guaiacolate).

#### SPECIAL PRECAUTIONS FOR THE OPERATOR

---

- Guaifenesin (Glyceryl Guaiacolate) is irritant to skin and mucous membranes, for that reason people should avoid any direct contact or accidental ingestion. The use of disposable gloves is recommended when handling this product.
- Wash hands after using the product.
- Guaifenesin (Glyceryl Guaiacolate) may cause hypersensitivity reactions.

#### ANTIDOTE

---

There is not specific antidote.

#### WARNINGS

---

Keep out of reach of children.

#### ADVERSE EFFECTS

---

Symptoms of breathing apnea, nystagmus, hypotension and stiff muscles are associated with toxic drug levels.

One liter of the product can be administered without showing toxic effects in the individual.

#### GUARD PERIOD

---

Do not use in horses which meat is intended for human consumption.

#### CONSERVATION

---

Store in a cool and dry place, between 15° and 30°C, protected from sunlight.

#### CONDITION OF SALE

---

To be supplied only on veterinary prescription.

#### PRESENTATION

---

serum container with 500 mL

#### PREPARED BY

---

Drag Pharma Laboratory.

#### RECORDS

---

Reg. SAG No. 2039  
Costa Rica: Reg. N° MAG CL 4-3-11-6176  
Perú: Reg. SENASA F.A0.01.I.0004

#### COUNTRIES WHERE IT IS MARKETED

---

Imported and Distributed by Representaciones Durand SAC.  
Av. Manuel Olgún No. 501 Office No. 604 Santiago de Surco Lima.

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian