

# INVECTINA® Injectable solution

## ENDECTOCIDE BROAD SPECTRUM ANTIPARASITIC



Endectocide indicated in the treatment of gastrointestinal parasites (*Ostertagia sp.*, *Nematodirus sp.*, *Oesophagostomum sp.*, *Haemonchus sp.*, *Cooperia sp.*, *Strongylus sp.*, *Trichuris sp.*, *Ascaris sp.*, *Trichostrongylus sp.*) And lung (*Dictyocaulus sp.*), including their migratory and inhibited larvae. Effective in treating scabies mites (*Sarcoptes scabiei*, *Psoroptes sp.*), Lice (*Linognathus sp.*, *Haematopinus sp.*) And *Oestrus ovis*.

## Technical Specification

### SPECIES

Bovine, sheep, goat and pigs.

### DOSAGE FORM

Injectable solution.

### THERAPEUTIC ACTION

Endectocide Broad spectrum Antiparasitic

### COMPOSITION

Each 1 mL of injectable solution contains:

Ivermectin.....10 mg  
Excipients q.s.p.....1 mL

### INDICATIONS

Endectocide indicated in the treatment of gastrointestinal parasites (*Ostertagia sp.*, *Nematodirus sp.*, *Oesophagostomum sp.*, *Haemonchus sp.*, *Cooperia sp.*, *Strongylus sp.*, *Trichuris sp.*, *Ascaris sp.*, *Trichostrongylus sp.*) And lung (*Dictyocaulus sp.*), including their migratory and inhibited larvae. Effective in treating scabies mites (*Sarcoptes scabiei*, *Psoroptes sp.*), Lice (*Linognathus sp.*, *Haematopinus sp.*) And *Oestrus ovis*.

### ROUTE OF ADMINISTRATION AND DOSAGE

#### Administration route:

Subcutaneous

#### Dosage and method of use:

- Cattle, sheep and goats: 1 mL / 50 Kg of weight (equivalent to 0.2 mg of Ivermectin / Kg of

weight) in a single dose. Repeat the dose after 7 days in case of sheep scabies.

- Pigs: 3 mL / 100 Kg of weight (equivalent to 0.3 mg of Ivermectin / Kg of weight) in a single dose.

Do not administer more than 10 mL per injection point.

#### DRUG INTERACTIONS

---

It is not recommended to use concomitantly with other pharmaceuticals.

#### CONTRAINDICATIONS

---

- Do not administer to animals younger than 4 weeks.
- Do not use in animals hypersensitive to Ivermectin

#### SPECIAL PRECAUTIONS FOR THE OPERATOR

---

- Do not handle by people hypersensitive to Ivermectin
- In case of contact with skin, eyes or mucosa, wash immediately with plenty of water.
- Do not smoke, drink or eat while administering the product.
- In case of injection, immediately go to a medical center and show the product label.

#### WARNINGS

---

Keep out of reach of children.

#### ADVERSE EFFECTS

---

Inflammation may occur at the injection site that resolves without treatment.

#### GUARD PERIOD

---

**Meat:** 49 days.

**Milk:** Do not use milk from treated animals for human consumption.

#### OBSERVATIONS

---

##### **Special precautions for the disposal of unused product or waste material:**

Discard the remains of unused product in its original container. Do not throw the empty container or with product remains, in rivers, lakes or streams of water, since the product is extremely toxic to aquatic species. Do not throw empty containers or containers with the rest of the product together with household waste. Do not reuse the container.

#### CONSERVATION

---

Store between 15 and 30 ° C, protected from light. Once the container is opened, use within 12 weeks. Discard the unused product after that period of time.

#### CONDITION OF SALE

---

Sale under Veterinary Medical prescription.

#### PRESENTATION

---

Ampule bottle with 50 mL and 100 mL, bottle with 500 mL

#### PREPARED BY

---

Drag Pharma Laboratory.

#### RECORDS

---

**Chile:** Reg. SAG N° 623-B

**Bolivia:** Reg. SENASAG PUV-F N° 007257/16

**Perú:** Registro SENASA F.09.01.I.0239

#### COUNTRIES WHERE IT IS MARKETED

---

**Imported and Distributed in Bolivia by:**

AGROGUARANI SRL

TEL: +(591)314-1401

Santa Cruz de la Sierra-Bolivia

**Imported and Distributed in Peru by:**

Representaciones Durand SAC.

Av. Manuel Olguín N° 501 Oficina N° 604 Santiago de Surco Lima.

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian